

vaxxino

Vaxxon[®] EDS



VACCINATE AGAINST EGG DROP SYNDROME

USE FROM 14 TO 20 WEEKS FOR LAYERS AND BREEDER HENS

ACTIVE SUBSTANCES: Inactivated Egg Drop Syndrome '76

ADMINISTRATION ROUTE: Intramuscularly to the chest or subcutaneously in the lower part of the neck.

DOSE: 0.5 ml

ADJUVANT: Light mineral oil. Shake well before use.

INDICATIONS FOR USE, SPECIFYING THE TARGET SPECIES: Vaccinate the laying hens and breeders from 14 to 20 weeks of age, before the entry in lay to achieve a durable protection against Egg Drop Syndrome '76.

SPECIAL PRECAUTIONS:

- Store and transport in a refrigerator (2° C – 8° C).
- Do not freeze.
- Protect from light.
- Allow bottles to reach room temperature (+ 20° C) prior to inoculation.
- For administration, appropriate aseptic conditions should be maintained.



EDS: Since its initial description, EDS virus has become a major cause of lost egg production throughout the world. The disease is characterized by the production of thin-shelled or shell-less eggs by otherwise healthy birds.

Once established in a breeding organization, the condition more often is seen as a failure to achieve production targets, and eggshell changes are less apparent.

Since its initial recognition, it has become apparent that sporadic outbreaks of EDS occur as a result of fowl becoming infected through direct or indirect contact with infected wild or domestic waterfowl.

vaxxino

1801 Biotech Ave. NE Willmar, MN 56201 | +1 320.222.9822
info.us@vaxxino.com | www.vaxxino.us.com

This product is for emergency use only, in premises approved by the State Veterinarian, in states identified by APHIS as positive for Egg Drop Syndrome.